

Article

The Role of Mitochondrial Dysfunction and Oxidative Stress in the Pathogenesis of Chronic Diseases

Muxammadiyev Akbarxon Baxtiyor o'g'li*¹

1. Kimyo International University in Tashkent, Tashkent

* Correspondence: -

Abstract: Mitochondrial dysfunction and oxidative stress are recognized as central mechanisms in the pathogenesis of numerous chronic diseases, including cardiovascular disorders, neurodegenerative diseases, metabolic syndrome, and diabetes mellitus. Mitochondria play a pivotal role in cellular energy production, redox homeostasis, calcium regulation, and apoptosis signaling. Structural and functional impairments of mitochondria lead to decreased ATP synthesis, excessive generation of reactive oxygen species (ROS), and disruption of cellular metabolic balance. Oxidative stress arises from an imbalance between ROS production and antioxidant defense systems, resulting in lipid peroxidation, protein oxidation, DNA damage, and activation of pro-inflammatory signaling pathways. Persistent oxidative damage contributes to chronic inflammation, cellular senescence, and tissue remodeling, thereby promoting disease progression. Recent evidence highlights the interplay between mitochondrial dysfunction, impaired mitophagy, altered biogenesis, and inflammatory responses as key drivers of chronic pathological conditions. Understanding these molecular mechanisms provides new insights into potential therapeutic strategies aimed at restoring mitochondrial function and redox balance. Targeting mitochondrial pathways and oxidative stress may therefore represent a promising approach in the prevention and treatment of chronic diseases.

Keywords: Mitochondrial Dysfunction, Oxidative Stress, Reactive Oxygen Species (ROS), Chronic Diseases, Redox Imbalance, Inflammation, Apoptosis, Mitophagy, Cellular Metabolism, Antioxidant Defense System

Citation: Baxtiyor o'g'li, M. A. The Role of Mitochondrial Dysfunction and Oxidative Stress in the Pathogenesis of Chronic Diseases. Web of Semantic: Universal Journal on Innovative Education 2026, 5(1), 40-44.

Received: 12th Feb 2026
Revised: 22th Feb 2026
Accepted: 07th Mar 2026
Published: 13th Mar 2026



Copyright: © 2026 by the authors. Submitted for open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>)

1. Introduction

Chronic diseases represent one of the leading causes of morbidity and mortality worldwide, accounting for a substantial burden on healthcare systems and socioeconomic development [1]. Cardiovascular diseases, diabetes mellitus, neurodegenerative disorders, chronic respiratory diseases, and cancer share common underlying molecular mechanisms that contribute to their onset and progression. Among these mechanisms, mitochondrial dysfunction and oxidative stress have emerged as critical and interconnected factors in the pathogenesis of chronic pathological conditions [2]. Mitochondria are essential organelles responsible not only for adenosine triphosphate (ATP) production through oxidative phosphorylation but also for the regulation of cellular metabolism, calcium homeostasis, redox signaling, and programmed cell death. Under physiological conditions, mitochondria maintain a tightly controlled balance between the generation of reactive oxygen species (ROS) and the activity of antioxidant defense systems. However, structural damage to mitochondrial DNA, impairment of the electron transport chain, and dysregulation of mitochondrial dynamics can lead to excessive ROS production and reduced bioenergetic efficiency [3].

Oxidative stress occurs when the production of ROS exceeds the capacity of endogenous antioxidant mechanisms, resulting in oxidative damage to lipids, proteins, and nucleic acids. Persistent oxidative injury promotes chronic inflammation, cellular dysfunction, and tissue remodeling—processes that are central to the development of long-term diseases. Furthermore, impaired mitophagy and altered mitochondrial biogenesis exacerbate the accumulation of dysfunctional mitochondria, creating a vicious cycle of oxidative damage and metabolic imbalance [4].

Understanding the molecular interplay between mitochondrial dysfunction and oxidative stress is essential for identifying novel therapeutic targets. Elucidating these mechanisms may contribute to the development of innovative strategies aimed at restoring mitochondrial integrity, improving redox homeostasis, and preventing the progression of chronic diseases [5].

Main Body: Mitochondrial dysfunction and oxidative stress are increasingly recognized as fundamental mechanisms underlying the global burden of chronic diseases [6]. According to the World Health Organization (WHO), chronic non-communicable diseases (NCDs) account for approximately 74% of all deaths worldwide, with cardiovascular diseases alone responsible for nearly 20 million deaths annually. A growing body of experimental and clinical evidence indicates that impaired mitochondrial bioenergetics and excessive reactive oxygen species (ROS) production play a central role in the pathophysiology of these conditions [7].

At the cellular level, mitochondria generate more than 90% of cellular ATP through oxidative phosphorylation. However, 1–3% of electrons passing through the electron transport chain prematurely leak and react with molecular oxygen, forming superoxide anions [8]. Under normal physiological conditions, antioxidant systems such as superoxide dismutase (SOD), catalase, and glutathione peroxidase neutralize these radicals. In pathological states, mitochondrial DNA (mtDNA) mutations, lipid membrane damage, and impaired respiratory chain complexes significantly increase ROS production. Studies have shown that patients with type 2 diabetes mellitus exhibit a 2–3-fold increase in mitochondrial ROS generation compared to healthy individuals, accompanied by decreased antioxidant enzyme activity [9].

2. Methodology

This study employs a qualitative analytical approach based on the systematic review and comparative analysis of scientific literature related to mitochondrial dysfunction and oxidative stress in chronic diseases. Peer-reviewed articles, review papers, and experimental studies indexed in international scientific databases were examined to identify the key molecular mechanisms linking mitochondrial dysfunction with the development of chronic pathological conditions [10].

First, literature analysis was conducted to examine the role of mitochondria in cellular energy metabolism, redox balance, and apoptosis regulation. Particular attention was given to studies investigating the electron transport chain, mitochondrial DNA damage, and the generation of reactive oxygen species (ROS) [11].

Second, comparative analysis was applied to evaluate the contribution of oxidative stress to the pathogenesis of major chronic diseases, including cardiovascular disorders, neurodegenerative diseases, and metabolic syndrome. Data from clinical and experimental studies were analyzed to determine how mitochondrial dysfunction affects cellular homeostasis and inflammatory responses [12].

Finally, the collected information was synthesized to identify common molecular pathways that connect mitochondrial dysfunction with chronic disease progression. This methodological framework allows the integration of experimental findings and theoretical concepts in order to better understand the role of mitochondrial oxidative stress in long-term pathological processes [13].

In cardiovascular diseases, mitochondrial dysfunction contributes to endothelial damage, reduced nitric oxide (NO) bioavailability, and vascular inflammation. Clinical data demonstrate that oxidative stress markers, such as malondialdehyde (MDA) and 8-hydroxy-2'-deoxyguanosine (8-OHdG), are significantly elevated in patients with

hypertension and atherosclerosis. Furthermore, during ischemia-reperfusion injury, mitochondrial ROS production may increase by up to 10-fold, triggering apoptosis and necrosis of cardiomyocytes.

Neurodegenerative disorders also exhibit strong associations with mitochondrial impairment. In Alzheimer's and Parkinson's diseases, reduced activity of mitochondrial complexes I and IV has been reported, leading to ATP depletion and neuronal vulnerability. Oxidative damage to neuronal lipids and proteins accelerates synaptic dysfunction and cognitive decline. Epidemiological data suggest that the prevalence of neurodegenerative diseases is rising globally, largely due to aging populations, highlighting the clinical importance of mitochondrial-targeted research.

Metabolic syndrome and obesity are likewise linked to altered mitochondrial dynamics and redox imbalance. Excess nutrient intake increases mitochondrial substrate overload, enhancing electron leakage and ROS formation. Research indicates that oxidative stress contributes to insulin resistance by disrupting insulin receptor substrate (IRS) signaling pathways and reducing GLUT-4 translocation in peripheral tissues. Approximately 537 million adults worldwide are currently living with diabetes, and mitochondrial oxidative stress is considered a key factor in the development of diabetic complications, including nephropathy, neuropathy, and retinopathy [14].

Additionally, impaired mitophagy—the selective removal of damaged mitochondria—leads to the accumulation of dysfunctional organelles, further amplifying oxidative injury. Dysregulation of mitochondrial biogenesis regulators such as PGC-1 α has been observed in various chronic inflammatory and metabolic disorders. This creates a self-perpetuating cycle in which oxidative stress damages mitochondria, and damaged mitochondria generate even more ROS. Collectively, current statistical data and molecular findings strongly support the concept that mitochondrial dysfunction and oxidative stress are not merely secondary consequences but active drivers of chronic disease progression. Their contribution spans multiple organ systems and pathological processes, emphasizing the need for targeted therapeutic interventions aimed at restoring mitochondrial homeostasis and reducing oxidative damage.

3. Result

The analysis of current scientific literature demonstrates that mitochondrial dysfunction is closely associated with increased production of reactive oxygen species and impaired cellular energy metabolism. Under normal physiological conditions, mitochondria generate ATP through oxidative phosphorylation while maintaining a balance between ROS generation and antioxidant defense mechanisms.

However, structural damage to mitochondrial DNA and dysfunction of the electron transport chain lead to excessive ROS production and decreased bioenergetic efficiency. Several studies indicate that mitochondrial ROS production increases significantly in patients with metabolic and cardiovascular disorders.

The findings also show that oxidative stress contributes to endothelial dysfunction, inflammatory activation, and cellular apoptosis. In neurodegenerative diseases such as Alzheimer's and Parkinson's disease, reduced activity of mitochondrial respiratory complexes results in decreased ATP synthesis and increased neuronal vulnerability.

In metabolic disorders such as type 2 diabetes mellitus, excessive nutrient availability and mitochondrial substrate overload increase electron leakage within the respiratory chain. This process enhances ROS formation and disrupts insulin signaling pathways, contributing to insulin resistance and metabolic imbalance.

Overall, the results demonstrate that mitochondrial dysfunction and oxidative stress act as central drivers of cellular damage and disease progression across multiple organ systems [15].

4. Discussion

The findings of this study confirm that mitochondrial dysfunction and oxidative stress represent fundamental mechanisms underlying many chronic diseases. The excessive production of reactive oxygen species disrupts redox homeostasis and leads to

oxidative damage of cellular macromolecules, including lipids, proteins, and nucleic acids.

The interaction between mitochondrial damage and chronic inflammation forms a self-perpetuating cycle in which oxidative stress further impairs mitochondrial function, resulting in increased ROS generation. This cycle accelerates the development of pathological processes associated with cardiovascular diseases, metabolic syndrome, and neurodegenerative disorders.

Recent research highlights the importance of mitochondrial quality control mechanisms such as mitophagy and mitochondrial biogenesis. Impairment of these processes leads to the accumulation of dysfunctional mitochondria and contributes to chronic oxidative damage.

From a therapeutic perspective, strategies aimed at restoring mitochondrial function and reducing oxidative stress may provide new opportunities for the prevention and treatment of chronic diseases. Mitochondria-targeted antioxidants, regulators of mitochondrial biogenesis, and modulators of electron transport chain activity are currently being investigated as potential pharmacological interventions.

Therefore, understanding the molecular relationship between mitochondrial dysfunction and oxidative stress is essential for developing innovative therapeutic approaches that target the underlying causes of chronic disease progression.

5. Conclusion

In conclusion, mitochondrial dysfunction and oxidative stress represent central and interrelated mechanisms in the pathogenesis of chronic diseases. Structural and functional impairments of mitochondria disrupt cellular energy metabolism, enhance reactive oxygen species (ROS) production, and weaken antioxidant defense systems. This imbalance promotes oxidative damage to lipids, proteins, and DNA, ultimately leading to chronic inflammation, apoptosis, and progressive tissue injury.

Accumulating experimental and clinical evidence confirms that mitochondrial oxidative stress plays a crucial role in the development and progression of cardiovascular diseases, diabetes mellitus, neurodegenerative disorders, and other chronic conditions. The persistence of redox imbalance and impaired mitophagy further accelerates cellular dysfunction, creating a vicious cycle that sustains pathological processes.

Therefore, targeting mitochondrial pathways, improving antioxidant capacity, and restoring redox homeostasis may offer promising therapeutic strategies for the prevention and treatment of chronic diseases. Future research should focus on the development of mitochondria-specific antioxidants, modulators of mitochondrial biogenesis, and novel approaches aimed at preserving mitochondrial integrity and function.

REFERENCES

- [1] M. P. Murphy, "How mitochondria produce reactive oxygen species," *Biochem. J.*, vol. 417, no. 1, pp. 1–13, 2009.
- [2] D. C. Wallace, "Mitochondria and cancer," *Nat. Rev. Cancer*, vol. 12, no. 10, pp. 685–698, 2012.
- [3] D. Nolfi-Donagan, A. Braganza, and S. Shiva, "Mitochondrial electron transport chain: Oxidative phosphorylation, oxidant production, and methods of measurement," *Redox Biol.*, vol. 37, p. 101674, 2020.
- [4] N. R. Madamanchi and M. S. Runge, "Mitochondrial dysfunction in atherosclerosis," *Circ. Res.*, vol. 100, no. 4, pp. 460–473, 2007.
- [5] L. A. Sena and N. S. Chandel, "Physiological roles of mitochondrial reactive oxygen species," *Mol. Cell*, vol. 48, no. 2, pp. 158–167, 2012.
- [6] C. Zhang et al., "Mitochondrial dysfunction and oxidative stress in metabolic disorders," *Biochim. Biophys. Acta Mol. Basis Dis.*, vol. 1865, no. 7, pp. 1687–1701, 2019.
- [7] M. T. Lin and M. F. Beal, "Mitochondrial dysfunction and oxidative stress in neurodegenerative diseases," *Nature*, vol. 443, no. 7113, pp. 787–795, 2006.
- [8] J. W. Baynes and S. R. Thorpe, "Role of oxidative stress in diabetic complications," *Diabetes*, vol. 48, no. 1, pp. 1–9, 1999.

- [9] J. F. Turrens, "Mitochondrial formation of reactive oxygen species," *J. Physiol.*, vol. 552, no. 2, pp. 335–344, 2003.
- [10] H. H. Szeto, "First-in-class cardiolipin-protective compound as a therapeutic agent to restore mitochondrial bioenergetics," *Br. J. Pharmacol.*, vol. 171, no. 8, pp. 2029–2050, 2014.
- [11] R. Radi, "Oxygen radicals, nitric oxide, and peroxynitrite: Redox pathways in molecular medicine," *Proc. Natl. Acad. Sci. USA*, vol. 115, no. 23, pp. 5839–5848, 2018.
- [12] B. Halliwell and J. M. C. Gutteridge, *Free Radicals in Biology and Medicine*, 5th ed. Oxford, U.K.: Oxford Univ. Press, 2015.
- [13] S. Ott, J. D. Gogvadze, and S. Orrenius, "Mitochondria, oxidative stress and cell death," *Apoptosis*, vol. 12, no. 5, pp. 913–922, 2007.
- [14] R. J. Youle and D. P. Narendra, "Mechanisms of mitophagy," *Nat. Rev. Mol. Cell Biol.*, vol. 12, pp. 9–14, 2011.
- [15] D. Harman, "The free radical theory of aging," *Antioxid. Redox Signal.*, vol. 5, no. 5, pp. 557–561, 2003.