

Integrated System of Measures for the Treatment of Papillomavirus Infection

Munisa Khamrakulovna Ablakulova, Abduxamidova Dilshoda Khalimovna

General Practice of Family Medicine, Samarkand State Medical University, Samarkand,
Uzbekistan

Article Information

Received: November 29, 2022

Accepted: December 30, 2022

Published: January 31, 2023

Keywords: Human papillomaviruses, treatment, disease recurrence.

ABSTRACT

this article attempts to reveal the main reasons for the creation of a comprehensive system of measures for the treatment of human papillomavirus infection (PVI) and a significant reduction in the recurrence of the disease. To carry out scientific work, the author found that HPV type 16 among patients with cervical cancer is detected in 68.3% of cases, HPV type 18 - in 11.7% of cases. A comparative evaluation of the effectiveness of monotherapy and combined treatment of PVI was carried out in 98 patients (age 25–37 years) with localization of tumor elements in the external genital area. Group I (52 people) consisted of patients who were treated only with the method of laser therapy (copper vapor laser 578 nm "Yakhroma-Med", which selectively affects the pathological tissue). Group II (46 people) consisted of patients who received combined treatment: laser destruction, epithelialization occurred within 5–7 days, after which Keravort cream (5% imiquimod) was administered externally, which was applied to the entire surface of the skin where condylomas were localized, capturing visibly healthy skin 1 cm in diameter at night 3 times a week for 4–8 weeks. The problem in question is still little studied, therefore, requires more thorough research.

Introduction: Human papillomaviruses (HPV), or human papillomaviruses (HPV, English human papillomavirus, HPV) are a group of viruses from the papillomavirus family, including 27 species from 5 genera (Alphapapillomavirus, Betapapillomavirus, Gammapapillomavirus, Mupapillomavirus and Nupapillomavirus) and more than 170 types (strains).

About 80% of the sexually active population becomes infected with HPV during their lifetime. 660 million people in the world are already infected with HPV (12% of the world's population). 750 thousand cases of HPV-associated cancers and 32 million cases of anogenital warts are registered annually in the world. Every year, more than 300 thousand people die from cancer caused by HPV[7][8][9]. HPV is the most common sexually transmitted disease in the United States, Russia and many other countries.

In itself, HPV infection does not mean evidence of sexual contact, since the standard life cycle of the virus is designed for infection through microcracks in the skin. Infection through the mucous membranes is only more effective and therefore more common. Infection with the virus is possible through shared objects or even a handshake[11][12]. Infection through objects and the skin of other people contributes to the extreme resistance of HPV to antiseptics.

HPV refers to non-enveloped viruses, that is, with a simple, but highly resistant to antiseptics, icosahedral capsid. Inside the capsid, the virus contains double-stranded DNA in the form of a

closed ring. The DNA of the virus duplicates its proteins many times, since the life cycle of the virus requires the breaking off of parts of the DNA[24]

Purpose: to create a comprehensive system of measures for the treatment of human papillomavirus infection (PVI) and a significant reduction in the recurrence of the disease.

Materials and methods: it was found that HPV type 16 among patients with cervical cancer is detected in 68.3% of cases, HPV type 18 - in 11.7% of cases. A comparative evaluation of the effectiveness of monotherapy and combined treatment of PVI was carried out in 98 patients (age 25–37 years) with localization of tumor elements in the external genital area. Group I (52 people) consisted of patients who were treated only with the method of laser therapy (copper vapor laser 578 nm "Yakhroma-Med", which selectively affects the pathological tissue). Group II (46 people) consisted of patients who received combined treatment: laser destruction, epithelialization occurred within 5–7 days, after which Keravort cream (5% imiquimod) was administered externally, which was applied to the entire surface of the skin where condylomas were localized, capturing visibly healthy skin 1 cm in diameter at night 3 times a week for 4-8 weeks. To assess the effectiveness of therapy, the following approaches were used: comparing the results of an HPV examination before the start of treatment and 4 months after the end of the course of treatment by PCR; assessment of the timing of resolution of clinical manifestations of PVI in patients receiving different treatment regimens; analysis of the frequency of relapses after treatment.

Results: the results of comparing the effectiveness of various PVI therapy options according to the criterion "relapse rate after treatment" turned out to be the most significant. In the first group, the recurrence rate was 18%, in the second group of patients, no relapses were noted.

Conclusions: Thus, the optimal tactic for the treatment of PVI is combined therapy, which includes the elimination of papillomatous foci by laser therapy using the Yakhroma-Med apparatus and local immunocorrection. At the same time, it should be noted that the drug "Keravort" is a means of etiotropic therapy for diseases associated with the human papillomavirus.

Literature:

1. Хусинова Ш., Аблакулова М., Хакимова Л. Процесс рационального назначения лекарственных средств и выбор персонального лекарства в практике ВОП //Общество и инновации. – 2020. – Т. 1. – №. 1/С. – С. 683-689.
2. Лапасов С. Х. и др. Результаты оценки качества диагностики и лечения больных сахарным диабетом 2 типа в условиях первичного звена медико-санитарной помощи //Здобутки клінічної і експериментальної медицини. – 2017. – №. 2. – С. 134-137.
3. Хакимова Л. и др. Безопасное материнство и эффективный антенатальный уход //Общество и инновации. – 2021. – Т. 2. – №. 8/С. – С. 91-95.
4. Хакимова Л. Р. и др. Результаты внедрения клинического протокола по интегрированному ведению больных с артериальной гипертензией и сахарным диабетом в первичном звене здравоохранения //Здоровье, демография, экология финно-угорских народов. – 2018. – №. 4. – С. 66-68.
5. Лапасов С. Х., Хусинова Ш. А., Хакимова Л. Р. Повышение качества информированности беременных женщин о подготовке к родам и методам контрацепции //Актуальные вопросы современной медицинской науки и здравоохранения: сборник статей IV Международной научно-практической конференции молодых учёных и студентов, IV Всероссийского форума медицинских и фармацевтических вузов «За качественное образование», (Екатеринбург, 10-12 апреля 2019): в 3-х т.-Екатеринбург: УГМУ, CD-

- ROM. – Федеральное государственное бюджетное образовательное учреждение высшего образования «Уральский государственный медицинский университет» Министерства здравоохранения Российской Федерации, 2019.
6. Лапасов С. Х. и др. Диагностика, лечение и профилактика хронического гепатита с в условиях первичного звена медицинской помощи (обзор литературы) //Здобутки клінічної і експериментальної медицини. – 2017. – №. 2. – С. 13-21.
 7. Хакимова Л. Р. и др. Оценка эффективности обучения основам доказательной медицины в последипломном образовании врачей общей практики //Медицина и экология. – 2017. – №. 4 (85). – С. 130-133.
 8. Рахимова Х. М., Сулаймонова Н. Э. Оптимизация методов лечения дисметаболического хронического пиелонефрита у детей //Здоровье, демография, экология финно-угорских народов. – 2019. – №. 1. – С. 54-56.
 9. Kholboyev S., Rahimova K., Sulaymanova N. The State of the Intestinal Microflora and Assessment of the Health State among People with Impaired Glucose Tolerance //Annals of the Romanian Society for Cell Biology. – 2021. – С. 163-176.
 10. Холбаев С. Б., Сулейманова Н. Э., Юлдашова Н. Э. Адаптированный клинический протокол по интегрированному ведению артериальной гипертензии и сахарного диабета среди лиц старше 40 лет //Вопросы неотложной кардиологии 2016. – 2016. – С. 6-7.
 11. Сулайманова Н. Олам лисоний манзарасининг баҳоли шаклланиши //Иностранная филология: язык, литература, образование. – 2021. – №. 2 (79). – С. 18-24.
 12. Сулайманова Н., Рахимова Х., Юлдашова Н. KEKSA YOSHDAGI BEMORLARGA DORI-DARMONLARNI TAYINLASH XUSUSIYATLARI //Журнал кардиореспираторных исследований. – 2021. – Т. 2. – №. 3. – С. 85-88.
 13. Kholbayev S. B. et al. THE EFFECT OF DRUG THERAPY IN PATIENTS WITH OBESITY ON THE CLINICAL MANIFESTATIONS OF KNEE OSTEOARTHRITIS. – 2020.
 14. Kholbayev S. B. et al. ВЛИЯНИЕ МЕДИКАМЕНТОЗНОЙ ТЕРАПИИ ОЖИРЕНИЯ НА КЛИНИЧЕСКИЕ ПРОЯВЛЕНИЯ ОСТЕОАРТРОЗА КОЛЕННЫХ СУСТАВОВ НА ПЕРВИЧНОМ ЗВЕНЕ ЗДРАВООХРАНЕНИЯ //Здобутки клінічної і експериментальної медицини. – №. 1. – С. 181-187.
 15. Холбоев С., Юлдашова Н. РЕЗУЛЬТАТЫ АНАЛИЗА ОСНОВАННЫЕ НА МИССИИ ВОЗ И КИТАЙСКОЙ НАРОДНОЙ РЕСПУБЛИКИ ПО ПРОБЛЕМЕ НОВОЙ КОРОНАВИРУСНОЙ ИНФЕКЦИИ COVID-2019 //Журнал кардиореспираторных исследований. – 2020. – Т. 1. – №. SI-1. – С. 81-81.
 16. Рахимова Х., Сулаймонова Н. AXOLI O'RTASIDA 60 VA UNDAN KATTA YOSHDAGI KISHILAR UCHUN KORONAVIRUS INFEKTSIYASINING OLDINI OLISH BO'YICHA TAVSIYALAR //Журнал кардиореспираторных исследований. – 2020. – Т. 1. – №. SI-1. – С. 85-85.
 17. Сулайманова Н. Э., Рахимова Х. М. YURAK QON TOMIR KASALLIKLARI ASORATLARI PROFILAKTASIDA KO 'RSATILADIGAN TIBBIY XIZMAT SIFATINI BAHOLASH //Журнал кардиореспираторных исследований. – 2022. – Т. 3. – №. 3.
 18. Rakhimova K. M., Soleeva S. S. Improving the quality of type 2 diabetes control based on pen protocols among the population //ACADEMICIA: An International Multidisciplinary Research Journal. – 2022. – Т. 12. – №. 3. – С. 62-68.

19. Sulaimanova N. E., Rakhimova K. M., Soleeva S. S. Polypharmacy in prescribing drugs to elderly patients //ACADEMICIA: An International Multidisciplinary Research Journal. – 2022. – Т. 12. – №. 3. – С. 55-61.
20. Akbarovna K. S. et al. REVIEW OF THE COURSE AND TREATMENT FEATURES OF COVID-19 PATIENTS WITH CONCOMITANT CARDIOVASCULAR DISEASE //Asian journal of pharmaceutical and biological research. – 2022. – Т. 11. – №. 2.
21. Рахимова Х., Сулаймонова Н. AXOLI O'RTASIDA 60 VA UN DAN KATTA YOSHDAGI KISHILAR UCHUN KORONAVIRUS INFEKTSIYASINING OLDINI OLISH BO'YICHA TAVSIYALAR //Журнал кардиореспираторных исследований. – 2020. – Т. 1. – №. SI-1. – С. 85-85.
22. Маматкулов Х. и др. Значение социальных факторов в психоэмоциональном развитии и формировании как личности детей школьного возраста от 10 до 14 лет //Журнал вестник врача. – 2011. – Т. 1. – №. 04. – С. 76-77.
23. Ахмеджанова Н., Эгамбердиева М., Рахимова Х. Влияние комплексного лечения на показатели эндогенной Интоксикации при хроническом пиелонефрите у детей //Журнал проблемы биологии и медицины. – 2018. – №. 1 (99). – С. 14-19.
24. Лапасов С. Х. и др. Инновационные подходы в диагностике язвенной болезни у взрослых в первичном звене здравоохранения: обзор литературы //Здоровье, демография, экология финно-угорских народов. – 2018. – №. 4. – С. 68-72.
25. Юлдашова Н., Хакимова Л. РОЛЬ СЕМЕЙНОЙ МЕДИЦИНСКОЙ СЕСТРЫ В РАБОТЕ «ШКОЛЫ МАТЕРИНСТВА» //InterConf. – 2021. – С. 507-512.
26. Юлдашова Н. РАСПРОСТРАНЕННОСТЬ ФАКТОРОВ РИСКА, ВЛИЯЮЩИХ НА ПРОГНОЗ ЗАБОЛЕВАНИЯ БОЛЬНЫХ С АРТЕРИАЛЬНОЙ ГИПЕРТЕНЗИЕЙ //InterConf. – 2020.
27. KHOLBOEV S. B., YULDASHOVA N. E. ADVANTAGES AND DISADVANTAGES OF ONLINE TRAINING SYSTEM WITH LEARNERS' POSITIONS //БИОМЕДИЦИНА ВА АМАЛИЁТ ЖУРНАЛИ. – С. 233.
28. Khakimova L. et al. Acne in Allergic Skin Diseases //Texas Journal of Medical Science. – 2022. – Т. 8. – С. 129-131.
29. Akbarovna K. S. et al. ASSESSMENT OF THE PREVALENCE AND QUALITY OF CARE OF PATIENTS WITH HEART FAILURE IN PRIMARY CARE //Thematics Journal of Education. – 2022. – Т. 7. – №. 3.
30. Аблакулова М., Абдухамидова Д. ОСОБЕННОСТИ СТАРЕЮЩЕГО ОРГАНИЗМА И ФАРМАКОТЕРАПИИ В ГЕРИАТРИИ //InterConf. – 2020.
31. Khakimova L. et al. acne in allergic skin diseases. texas journal of medical science, 8. – 1818.
32. Khakimova L. et al. Acne in Allergic Skin Diseases //Texas Journal of Medical Science. – 2022. – Т. 8. – С. 129-131.